

Local pressure for inhomogeneous fluids

James Dufty¹  | Jeffrey Wrighton¹  | Kai Luo² 

¹Department of Physics, University of Florida, Gainesville, Florida

²Department of Applied Physics, Nanjing University of Science and Technology, Nanjing, Jiangsu, 210094

Correspondence

James Dufty, Department of Physics,
University of Florida, Gainesville, FL 32611.
Email: dufty@phys.ufl.edu

Funding information

Carnegie Institution for Science.; National
Science Foundation, Grant/Award Number:
NSF CSEDI grant EAR-1901813; US
Department of Energy, Grant/Award Number:
Grant DE-SC0002139.

Abstract

Definitions for a local pressure in an inhomogeneous fluid are considered for both equilibrium and local equilibrium states. Thermodynamic and mechanical (hydrodynamic) contexts are reconciled. Remaining problems and uncertainties are discussed.

1 | INTRODUCTION

The global pressure for equilibrium systems is well defined within statistical mechanics, both quantum and classical.¹ For extensive systems, the definitions for different ensembles are equivalent. Here the grand canonical ensemble will be chosen for such representations, characterized by an inverse temperature β and activity ν . The global pressure is then proportional to the grand potential, which defines all thermodynamic properties of the system. Its form is obtained from variations of the grand potential with respect to volume, leading to the equilibrium average of a specific operator equivalent to the familiar virial equation. For the special case of inhomogeneous equilibrium states for systems with an external potential $v^{\text{ext}}(\mathbf{r})$ the effective activity $\nu(\mathbf{r}) \equiv \nu - \beta v^{\text{ext}}(\mathbf{r})$ varies locally so it is appropriate to define an associated local thermodynamics.² The first objective here is to explore how to define a local thermodynamic pressure $p^e(\mathbf{r}, \beta | \nu)$ whose spatial integral is the global pressure.³⁻⁵ The notation indicates that the local pressure depends on the space point \mathbf{r} , the inverse temperature β , and is a functional of $\nu(\mathbf{r})$. It is straightforward to identify a local operator whose ensemble average integrates to the global virial equation, defining such a local pressure. However, it is not unique since any contribution whose integral vanishes can be added. Additional constraints are needed.

A conceptually different notion of pressure is obtained from the average momentum flux of the inhomogeneous equilibrium fluid, or pressure tensor $p_{ij}^e(\mathbf{r}, \beta | \nu)$. Local conservation of momentum at equilibrium leads to a force balance equation relating the gradient of that pressure tensor to the applied external force. This approach has an

extensive history in the classical description of interfaces and surface tension.⁶⁻⁹ Derivation of the conservation law from the underlying Heisenberg dynamics provides the form of the operator whose average gives the pressure tensor. It does not explicitly exploit the grand potential or any thermodynamics other than the stationarity of the equilibrium state. While the local thermodynamic pressure is defined only indirectly from the global pressure, the pressure tensor is inherently a local property. However, this method only provides the divergence of the pressure tensor and the latter is therefore not unique. Consequently the related scalar pressure $p_m^e(\mathbf{r}, \beta | \nu) \equiv p_{ii}^e(\mathbf{r}, \beta | \nu)/3$ also is not unique (here and below a summation over repeated indices is implied). The pressure obtained from the pressure tensor will be referred to as the hydrodynamic or mechanical pressure as it appears in the macroscopic conservation equations. Clearly, it is desirable that the thermodynamic and hydrodynamic pressures should be the same for consistent representations of the stationary states. It is expected that the uncertainties in each can be exploited to assure this equivalence.

Two cases are considered here, the inhomogeneous equilibrium states described above, and their generalization to local equilibrium states. The latter differ in the sense that the inverse temperature can be spatially varying, $\beta = \beta(\mathbf{r})$, in addition to the activity $\nu(\mathbf{r})$. For equilibrium states it is shown that the uncertainty in the thermodynamic pressure can be removed by adding a contribution that equates it to the hydrodynamic pressure.

The same objective arises in the more general context of non-stationary local equilibrium states of hydrodynamics. The associated ensemble is similar to the equilibrium ensemble. A “thermodynamics”

for this state can be defined from the associated grand potential and an associated local pressure identified.^{10,11} However, in this case, for spatially varying $\beta(\mathbf{r})$, there is no longer the flexibility to modify the thermodynamic local pressure to be equal to that from the local equilibrium average of the stress tensor. Consequently, it would seem that the equation of state for hydrodynamics is not the same as that for local equilibrium thermodynamics. The precise difference is identified below. Unfortunately, this implies that the equivalence chosen for the strict equilibrium noted above is not recovered from the hydrodynamic equations for their stationary limit.

This paradox is resolved by exploiting the uncertainty in the pressure tensor. A divergenceless additional contribution to the average momentum flux can be chosen such that the local pressure associated with the new momentum flux agrees with the thermodynamic pressure. In this way, thermodynamic and mechanical concepts are reconciled.

The analysis here is based in quantum statistical mechanics so that all average properties have an associated underlying operator representing them. The calculation of their averages is not discussed but the connection to density functional theory methods is indicated. An alternative approach is to postulate an average pressure tensor and verify that it yields the required macroscopic force balance equation. This has been described by Percus⁵ for the inhomogeneous equilibrium fluid. His pressure tensor is entirely characterized by the thermodynamic free energy density. It is described in Supporting Information S3 and the associated pressure is identified.

The primary importance of this investigation of equivalence is for the local equilibrium states of hydrodynamics. In that case both concepts of the pressure occur. The first is as a functional relationship between the fundamental conserved number, energy, and momentum densities and their conjugate fields activity, temperature, and flow velocity. This functional relationship is the thermodynamics of the local equilibrium grand potential, or thermodynamic pressure. The second occurrence is through the average momentum flux, comprised as a reference local equilibrium average and a dissipative component. Only the reference contribution is considered here and is referred to as the average mechanical pressure tensor. It is a functional of the conjugate fields. Hence the equivalence of the thermodynamic and mechanical pressures is a necessary condition for the hydrodynamic equations to provide a closed local macroscopic description, regardless of the choice for the dissipative component (e.g., Navier–Stokes or far from equilibrium).

At this point it is appropriate to characterize the context by noting related topics not bearing directly on the question of equivalence. The lack of uniqueness for the mechanical pressure tensor is well known; for early references see References 6–9. Most of these studies do not make explicit the required equivalence with thermodynamic pressure. An exception is one demonstration that the Harasima choice gives the wrong pressure in cylindrical coordinates.¹² This is resolved in a recent work for cylindrical geometry by synthesizing the Irving and Kirkwood and Harasima expressions for different coordinates.¹³ In Reference 14, another definition of the local thermodynamic pressure closer to that given here does not make any connection to the

various choices for the mechanical pressure tensor. In summary, the work here is complementary to this important body of work by relating the two different studies of thermodynamic pressure and mechanical pressure tensor. Further comment is given in the final section where measurement by simulation is briefly discussed.

The local equilibrium thermodynamics considered here describes a reference state for real nonequilibrium systems. In the context of information entropy^{15–17} it provides the optimal representation in terms of the given exact average local conserved densities. The latter must be provided from some detailed exact theory (e.g., Liouville–von Neumann equation). The grand potential associated with the local equilibrium ensemble provides the “equation of state” for generating conjugate variables such as temperature and activity, and is the direct analog of equilibrium thermodynamics. To further clarify the context it is noted that the current work does not relate to the general fields of “nonequilibrium thermodynamics”¹⁸ or “extended thermodynamics.”¹⁹ The former is an attempt to discover universal fundamental principles, similar to those of equilibrium thermodynamics (e.g., a generalized second law, entropy), that govern the dissipative dynamics of macroscopic properties. Extended irreversible thermodynamics takes as the macroscopic fields the usual local conserved fields plus the dissipative fluxes of energy and momentum. The conservation laws of ordinary hydrodynamics must then be supplemented with unknown additional equations for the dissipative fluxes. If the latter could be given in terms of the conserved fields this would not be necessary, so in a sense extended hydrodynamics is a tool to discover those forms. The local equilibrium thermodynamics considered here is not a theory, such as those sought above, but rather an exact functional relationship among two equivalence classes of fields—there is no entropy production nor inherent dissipation beyond that of the input fields. While it represents general nonequilibrium states, it is not predictive without the hydrodynamic equations themselves (for an exact formulation of the latter see Reference 10).

2 | LOCAL PRESSURE FOR AN INHOMOGENEOUS FLUID AT EQUILIBRIUM

Consider first a system of N particles in a large volume V with Hamiltonian

$$\mathcal{H}_N = H_N + \sum_{\alpha=1}^N v^{\text{ext}}(\mathbf{q}_{\alpha}), \quad (1)$$

where $v^{\text{ext}}(\mathbf{q}_{\alpha})$ is an external potential coupling to the particle with position operator \mathbf{q}_{α} , and the isolated system Hamiltonian H_N is

$$H_N = \sum_{\alpha=1}^N \frac{p_{\alpha}^2}{2m} + \frac{1}{2} \sum_{\alpha \neq \sigma=1}^N U_N(|\mathbf{q}_{\alpha} - \mathbf{q}_{\sigma}|). \quad (2)$$

Here $U_N(|\mathbf{q}_{\alpha} - \mathbf{q}_{\sigma}|)$ is a pair potential for particles α and σ , and \mathbf{p}_{α} is the momentum operator for particle α . At equilibrium with inverse

temperature β and activity ν the average of a property characterized by an operator X_N is given in the grand canonical ensemble by

$$\langle X \rangle^e \equiv \sum_N \text{Tr}^{(N)} X_N \rho_N^e, \quad \rho_N^e = e^{-Q^e(\beta, V|\nu)} e^{-(\beta H_N - \int d\mathbf{r} \nu(\mathbf{r}) n(\mathbf{r}))}. \quad (3)$$

The external potential has been combined with the activity to define a local activity $\nu(\mathbf{r})$

$$\nu(\mathbf{r}) \equiv \nu - \beta v^{\text{ext}}(\mathbf{r}), \quad (4)$$

and $n(\mathbf{r})$ is the number density operator

$$n(\mathbf{r}) = \sum_{\alpha=1}^N \delta(\mathbf{r} - \mathbf{q}_\alpha). \quad (5)$$

The notation $Y(\beta, V|\nu)$ here and below denotes a function of β , V and a functional of $\nu(\mathbf{r})$. The normalization function $Q^e(\beta, V|\nu)$ is known as the grand potential²⁰ and is chosen such that $\langle \mathbf{1} \rangle^e = 1$

$$Q^e(\beta, V|\nu) = \ln \sum_N \text{Tr}^{(N)} e^{-(\beta H_N - \int d\mathbf{r} \nu(\mathbf{r}) n(\mathbf{r}))}. \quad (6)$$

It determines the complete thermodynamics for the system through the definition of the global pressure

$$\beta P^e(\beta|\nu) V = Q^e(\beta, V|\nu). \quad (7)$$

For a sufficiently large volume $Q^e(\beta, V|\nu)$ is extensive (proportional to V) so the pressure is independent of the volume. Then an equivalent form for Equation (7) is

$$\beta P^e(\beta|\nu) = \left. \frac{\partial Q^e(\beta, V|\nu)}{\partial V} \right|_{\beta, \nu}. \quad (8)$$

The volume derivative can be calculated directly (e.g., using length scaling^{10,21}) to get

$$P^e(\beta|\nu) = \frac{1}{3V} (2\langle K \rangle^e + \langle \mathcal{V} \rangle^e), \quad (9)$$

where K is the kinetic energy operator and \mathcal{V} is the virial operator (for the internal forces)

$$K_N = \sum_{\alpha=1}^N \frac{1}{2m} p_{\alpha j}^2, \quad \mathcal{V}_N = \frac{1}{2} \sum_{\alpha \neq \gamma=1}^N (\mathbf{q}_\gamma - \mathbf{q}_\alpha) \cdot \mathbf{F}_{\alpha\gamma}(|\mathbf{q}_\alpha - \mathbf{q}_\gamma|). \quad (10)$$

It is seen that Equation (9) is the usual virial equation for the global (intrinsic) pressure, confirming the consistency of the thermodynamic and mechanical concepts of global pressure.

The objective now is to identify an associated local density pressure. It is done by defining a local density for the grand potential in Equation (7)

$$\int d\mathbf{r} \beta p^e(\mathbf{r}, \beta|\nu) = Q^e(\beta, V|\nu). \quad (11)$$

Accordingly, $V^{-1} p^e(\mathbf{r}, \beta|\nu)$ is the local density for the global pressure

$$P^e(\beta|\nu) \equiv \frac{1}{V} \int d\mathbf{r} p^e(\mathbf{r}, \beta|\nu). \quad (12)$$

One choice to identify it is to replace the operators K and \mathcal{V} in Equation (9) by associated local densities

$$p_0^e(\mathbf{r}, \beta|\nu) = \frac{1}{3} (2K_0(\mathbf{r}) + \mathcal{V}_0(\mathbf{r}))^e, \quad (13)$$

where

$$K_{0N}(\mathbf{r}) = \frac{1}{4m} \sum_{\alpha=1}^N [p_{\alpha}^2, \delta(\mathbf{r} - \mathbf{q}_\alpha)]_+, \quad (14)$$

$$\mathcal{V}_{0N}(\mathbf{r}) = \frac{1}{2} \sum_{\alpha \neq \sigma=1}^N F_{\alpha\sigma}(|\mathbf{q}_\sigma - \mathbf{q}_\alpha|) (\mathbf{q}_\sigma - \mathbf{q}_\alpha) \delta(\mathbf{r} - \mathbf{q}_\alpha). \quad (15)$$

The brackets $[a, b]_+$ denote an anticommutator. It is required in Equation (14) to assure that $K_{0N}(\mathbf{r})$ is Hermitian. More generally, the local pressure can be expressed as

$$p^e(\mathbf{r}, \beta|\nu) = p_0^e(\mathbf{r}, \beta|\nu) + \Delta p_0^e(\mathbf{r}, \beta|\nu), \quad (16)$$

where $\Delta p_0^e(\mathbf{r}, \beta|\nu)$ is any functional whose volume integral vanishes.

$$\int d\mathbf{r} \Delta p^e(\mathbf{r}, \beta|\nu) = 0. \quad (17)$$

To suggest an alternative choice for $p^e(\mathbf{r}, \beta|\nu)$ consider the exact microscopic conservation law for the momentum density operator $p_i(\mathbf{r}, t)$ (see Supporting Information S1)

$$\partial_t p_i(\mathbf{r}, t) + \partial_j t_{ij}(\mathbf{r}, t) = -n(\mathbf{r}, t) \partial_i v^{\text{ext}}(\mathbf{r}, t), \quad (18)$$

where the momentum density operator is

$$\mathbf{p}(\mathbf{r}, t) = \frac{1}{2} \sum_{\alpha=1}^N [\mathbf{p}_\alpha(t), \delta(\mathbf{r} - \mathbf{q}_\alpha(t))]_+, \quad (19)$$

and the momentum flux is

$$t_{ij}(\mathbf{r}, t) = \frac{1}{4m} \sum_{\alpha=1}^N \left[p_{i\alpha}(t), [p_{j\alpha}(t), \delta(\mathbf{r} - \mathbf{q}_\alpha(t))] \right]_+ + \frac{1}{2} \sum_{\alpha \neq \sigma=1}^N F_{\alpha\sigma i}(|\mathbf{q}_\alpha(t) - \mathbf{q}_\sigma(t)|) \mathcal{D}_j(\mathbf{r}, \mathbf{q}_\alpha(t), \mathbf{q}_\sigma(t)). \quad (20)$$

The operator $\mathcal{D}_j(\mathbf{r}, \mathbf{q}_\alpha(t), \mathbf{q}_\sigma(t))$ is given by

$$\mathcal{D}_j(\mathbf{r}, \mathbf{q}_1, \mathbf{q}_2) \equiv \int_C d\lambda \frac{dx_j(\lambda)}{d\lambda} \delta(\mathbf{r} - \mathbf{x}(\lambda)), \quad \mathbf{x}(\lambda_1) = \mathbf{q}_1, \quad \mathbf{x}(\lambda_2) = \mathbf{q}_2. \quad (21)$$

Here \mathcal{C} is an arbitrary continuous path connecting $\mathbf{x}(\lambda)$ between λ_1 and λ_2 . (Further comment on the choice is given in Supporting Information S2.)

Since the equilibrium ensemble is stationary, the equilibrium average of Equation (18) gives the stationary equilibrium force balance equation

$$\partial_j \langle t_{ij}(\mathbf{r}) \rangle^e = -\langle n(\mathbf{r}) \rangle^e \partial_i v^{\text{ext}}(\mathbf{r}). \quad (22)$$

This is the expected local stability condition. A mechanical pressure can be associated with the average momentum flux according to

$$p_m^e(\mathbf{r}, \beta|\nu) \equiv \frac{1}{3} \langle t_{ii}(\mathbf{r}) \rangle^e. \quad (23)$$

Note that Equation (20) implies

$$\frac{1}{V} \int d\mathbf{r} \frac{1}{3} \langle t_{ii}(\mathbf{r}) \rangle^e = \frac{1}{3V} (2\langle K \rangle^e + \langle \mathcal{V} \rangle^e) = P^e(\beta|\nu), \quad (24)$$

where the last equality follows from Equation (9). Use has been made of the identity

$$\int d\mathbf{r} \mathcal{D}(\mathbf{r}, \mathbf{q}_\alpha, \mathbf{q}_\gamma) = \mathbf{q}_\alpha - \mathbf{q}_\gamma. \quad (25)$$

The left side of Equation (24) suggests an alternative choice for the definition of a local pressure in Equation (12)

$$p^e(\mathbf{r}, \beta|\nu) = p_m^e(\mathbf{r}, \beta|\nu) = \frac{1}{3} \langle t_{ii}(\mathbf{r}) \rangle^e. \quad (26)$$

This definition has a mechanical origin, without direct reference to the grand potential or thermodynamics.

The two choices $p_0^e(\mathbf{r}, \beta|\nu)$ and $p_m^e(\mathbf{r}, \beta|\nu)$ are clearly different, but both yield the thermodynamic global pressure. The choice $p^e(\mathbf{r}, \beta|\nu) = p_m^e(\mathbf{r}, \beta|\nu)$ for the local pressure in Equation (12) is clearly the desirable one as it assures the mechanical balance equation is consistent with thermodynamics.

Since $t_{ij}(\mathbf{r})$ has a precise microscopic origin, this provides a local microscopic basis for the pressure

$$p^e(\mathbf{r}, \beta|\nu) = \frac{1}{3} (2\langle K(\mathbf{r}) \rangle^e + \langle \mathcal{V}(\mathbf{r}) \rangle^e), \quad (27)$$

with

$$K_N(\mathbf{r}) = \frac{1}{8m} \sum_{\alpha=1}^N [p_{i\alpha}, [p_{i\alpha}, \delta(\mathbf{r} - \mathbf{q}_\alpha)]]_+, \quad (28)$$

and

$$\mathcal{V}_N(\mathbf{r}) = \frac{1}{2} \sum_{\alpha \neq \sigma=1}^N F_{\alpha\sigma i}(|\mathbf{q}_\alpha - \mathbf{q}_\sigma|) \mathcal{D}_i(\mathbf{r}, \mathbf{q}_\alpha, \mathbf{q}_\sigma).$$

This is the desired result.

It is seen that while the global pressure is the same for $p^e(\mathbf{r}, \beta|\nu)$ and $p_0^e(\mathbf{r}, \beta|\nu)$, the local pressures differ in their microscopic realizations

$$K_N(\mathbf{r}) = K_{0N}(\mathbf{r}) + \frac{\hbar^2}{8m} \nabla^2 n(\mathbf{r}), \quad (29)$$

$$\mathcal{V}_N(\mathbf{r}) = \mathcal{V}_{0N}(\mathbf{r}) + \frac{1}{2} \sum_{\alpha \neq \sigma=1}^N F_{\alpha\sigma i}(|\mathbf{q}_\alpha - \mathbf{q}_\sigma|) [\mathcal{D}_i(\mathbf{r}, \mathbf{q}_\alpha, \mathbf{q}_\sigma) - (\mathbf{q}_\sigma - \mathbf{q}_\alpha)_i \delta(\mathbf{r} - \mathbf{q}_\alpha)]. \quad (30)$$

To get Equation (29) use has been made of the identity

$$\frac{1}{2m} \sum_{\alpha=1}^N p_{\alpha i} \delta(\mathbf{r} - \mathbf{q}_\alpha) p_{\alpha i} = K_{0N}(\mathbf{r}) + \frac{\hbar^2}{4m} \nabla^2 n(\mathbf{r}). \quad (31)$$

The dependence on \hbar has been made explicit in Equations (29) and (31) to emphasize this is a purely quantum effect. The identification of the local pressure in terms of the average momentum flux is a common definition. What is new here is its identification as the thermodynamic local pressure for the grand ensemble.

The local pressure $p_m^e(\mathbf{r}, \beta|\nu)$ is more sensitive to spatial variations of the inhomogeneous state than $p_0^e(\mathbf{r}, \beta|\nu)$. In fact $p_0^e(\mathbf{r}, \beta|\nu)$ results from it by a leading order Taylor series approximation, for example,

$$\begin{aligned} \mathcal{D}_i(\mathbf{r}, \mathbf{q}_\alpha, \mathbf{q}_\sigma) &= (\mathbf{q}_\sigma - \mathbf{q}_\alpha)_i \delta(\mathbf{r} - \mathbf{q}_\alpha) + \int_{\lambda_2}^{\lambda_1} d\lambda \frac{dx_i(\lambda)}{d\lambda} [\delta(\mathbf{r} - \mathbf{x}(\lambda)) - \delta(\mathbf{r} - \mathbf{q}_\alpha)] \\ &\simeq (\mathbf{q}_\sigma - \mathbf{q}_\alpha)_i \delta(\mathbf{r} - \mathbf{q}_\alpha). \end{aligned} \quad (32)$$

This gives $\mathcal{V}_N(\mathbf{r}) \simeq \mathcal{V}_{0N}(\mathbf{r})$ and $p_m^e(\mathbf{r}, \beta|\nu) \simeq p_0^e(\mathbf{r}, \beta|\nu)$. This leading order approximation is justified only in the context of states that have smooth spatial variations over distances of the order of the force range. For extreme conditions, such as occur for warm, dense matter states it would seem that the form Equation (20) must be used for the momentum flux.

In summary, two definitions for a local pressure have been identified, $p_0^e(\mathbf{r}, \beta|\nu)$ and $p_m^e(\mathbf{r}, \beta|\nu)$. Each has been identified in terms of the

average of an underlying microscopic operator. They provide the same global thermodynamics in the sense that both of their volume integrals yield $P^e(\beta|\nu)$. However, at the local level they differ by $\Delta p_0^e(\mathbf{r}, \beta|\nu)$, identified from Equations (29–31). The choice of $p^e(\mathbf{r}, \beta|\nu) = p_m^e(\mathbf{r}, \beta|\nu)$ for the thermodynamic pressure is made on the basis of equating thermodynamic and mechanical definitions.

So far only the scalar local pressure has been considered. There is no thermodynamic route to define a local pressure tensor for an inhomogeneous fluid at equilibrium (see, however, Supporting Information S3). Instead it is identified from Equation (22)

$$p_{ij}^e(\mathbf{r}, \beta|\nu) \equiv \langle t_{ij}(\mathbf{r}) \rangle^e. \quad (33)$$

By construction it has the form

$$p_{ij}^e(\mathbf{r}, \beta|\nu) = \frac{1}{3} p^e(\mathbf{r}, \beta|\nu) \delta_{ij} + \tilde{p}_{ij}^e(\mathbf{r}, \beta|\nu), \quad (34)$$

(where \tilde{p}_{ij}^e is its traceless part) and satisfies the force balance equation

$$\partial_i p_{ij}^e(\mathbf{r}, \beta|\nu) = -\langle n(\mathbf{r}) \rangle \partial_i v^{\text{ext}}(\mathbf{r}). \quad (35)$$

In the next section, attempts to extend the equivalence of thermodynamic and mechanical concepts to local equilibrium states in the same way lead to difficulties due to the spatial variations of $\beta(\mathbf{r}, t)$.

3 | LOCAL HYDRODYNAMIC PRESSURE

Consider now a general nonequilibrium state. The macroscopic hydrodynamic equations have their origins in averages of the underlying microscopic conservation laws for number density, energy density, and momentum density, $\{\langle n(\mathbf{r}, t) \rangle, \langle e(\mathbf{r}, t) \rangle, \langle \mathbf{p}(\mathbf{r}, t) \rangle\}$.^{10,11,22} In particular the hydrodynamic equation resulting from the conservation law for the momentum density follows from the nonequilibrium average of Equation (18)

$$\partial_t \langle \mathbf{p}(\mathbf{r}, t) \rangle + \partial_j \langle t_{ij}(\mathbf{r}, t) \rangle = -\langle n(\mathbf{r}, t) \rangle \partial_i v^{\text{ext}}(\mathbf{r}, t), \quad (36)$$

where the brackets now denote a nonequilibrium average

$$\langle X(t) \rangle \equiv \sum_N \text{Tr}^{(N)} X_N \rho_N(t), \quad (37)$$

and $\rho_N(t)$ is a solution to the Liouville–von Neumann equation. Traditionally, the momentum density is expressed in terms of a local flow velocity $\mathbf{u}(\mathbf{r}, t)$ defined by

$$\langle \mathbf{p}(\mathbf{r}, t) \rangle \equiv m \langle n(\mathbf{r}, t) \rangle \mathbf{u}(\mathbf{r}, t), \quad (38)$$

and Equation (36) is written in terms of the momentum flux in the local rest frame

$$\langle t_{ij}(\mathbf{r}, t) \rangle = m \langle n(\mathbf{r}, t) \rangle u_i(\mathbf{r}, t) u_j(\mathbf{r}, t) + \langle t_{0ij}(\mathbf{r}, t) \rangle. \quad (39)$$

Here the rest frame momentum flux $t_{0ij}(\mathbf{r}, t)$ has the same form as Equation (20) with the particle momenta in the rest frame, $p_{ia}(t) \rightarrow p_{ia}(t) - m u_i(\mathbf{q}_a(t), t)$. Then the momentum conservation law takes the form

$$D_t u_i(\mathbf{r}, t) + \partial_j \langle t_{0ij}(\mathbf{r}, t) \rangle = -\langle n(\mathbf{r}, t) \rangle \partial_i v^{\text{ext}}(\mathbf{r}, t), \quad (40)$$

with the material derivative $D_t = \partial_t + \mathbf{u}(\mathbf{r}, t) \cdot \nabla$. In this way, the purely convective contributions have been made explicit.

It remains to calculate the rest frame momentum flux $\langle t_{0ij}(\mathbf{r}, t) \rangle$. To do so, the solution to the Liouville–von Neumann equation is separated into a reference local equilibrium state, ρ_N^e , and its remainder Δ_N

$$\rho_N(t) = \rho_N^e[y(t)] + \Delta_N(t). \quad (41)$$

The reference local equilibrium state is chosen to be entirely determined by a set of conjugate fields $\{y(t)\}$ in one-to-one correspondence with the macroscopic conserved fields $\{\langle n(\mathbf{r}, t) \rangle, \langle e(\mathbf{r}, t) \rangle, \langle \mathbf{p}(\mathbf{r}, t) \rangle\}$. This correspondence is defined by the requirements that the reference state yield the exact averages for the local conserved fields

$$\bar{n}^e(\mathbf{r}|y(t)) \equiv \langle n(\mathbf{r}, t) \rangle, \quad (42a)$$

$$\bar{e}^e(\mathbf{r}|y(t)) \equiv \langle e(\mathbf{r}, t) \rangle, \quad (42b)$$

$$\bar{\mathbf{p}}^e(\mathbf{r}|y(t)) \equiv \langle \mathbf{p}(\mathbf{r}, t) \rangle, \quad (42c)$$

where the superscript e denotes a reference ensemble average, $\bar{A}^e = \langle A; \rho^e \rangle$. The left sides of these equations are functionals of the conjugate fields while the right sides are the fields of the local conservation laws. In this way, the conjugate fields $\{y(t)\}$ are functionals of the average conserved fields, and vice versa by inversion. The reference state therefore has the exact average values for the conserved fields by construction.

A choice for ρ_N^e with these properties is the local equilibrium ensemble^{11,22}

$$\rho_N^e[y(t)] = e^{-\eta[y(t)]}, \quad \eta[y(t)] = Q^e[y(t)] + \int d\mathbf{r} \psi_\kappa(\mathbf{r}) y_\kappa(\mathbf{r}, t), \quad (43)$$

$$Q^e[y(t)] = \ln \sum_N \text{Tr}^{(N)} e^{-\int d\mathbf{r} \psi_\kappa(\mathbf{r}) y_\kappa(\mathbf{r}, t)}, \quad (44)$$

where $\psi_\kappa(\mathbf{r})$ are the operators representing the local conserved number density, energy density, and momentum density

$$\{\psi_\kappa(\mathbf{r})\} \equiv \{n(\mathbf{r}), e(\mathbf{r}), \mathbf{p}(\mathbf{r})\}, \quad (45)$$

and $y_\kappa(\mathbf{r}, t)$ are the conjugate fields,

$$\{y(\mathbf{r}, t)\} \leftrightarrow \left\{ \left[-\nu(\mathbf{r}, t) + \frac{\beta(\mathbf{r}, t)}{2} m u^2(\mathbf{r}, t) \right], \beta(\mathbf{r}, t), -\beta(\mathbf{r}, t) \mathbf{u}(\mathbf{r}, t) \right\}. \quad (46)$$

It is interesting to note that this local equilibrium ensemble is also the “best choice” in the sense that it maximizes the information entropy for the given values of the conservative fields.¹⁵⁻¹⁷ In Reference 11 it is also called the “relevant” ensemble.

Accordingly, the average rest frame momentum flux $\langle t_{0ij}(\mathbf{r}, t) \rangle$ of Equation (39) has two contributions. One is from the reference local equilibrium ensemble ρ_N^ℓ and one from the remainder in Equation (41)

$$\begin{aligned} \langle t_{0ij}(\mathbf{r}, t) \rangle &= \sum_N Tr^{(N)} t_{0ij}(\mathbf{r}) \rho_N^\ell[y(t)] + \sum_N Tr^{(N)} t_{0ij}(\mathbf{r}) \Delta_N(t) \\ &\equiv \overline{t_{0ij}^\ell}(\mathbf{r}|y(t)) + \delta \langle t_{0ij}(\mathbf{r}, t) \rangle. \end{aligned} \quad (47)$$

It is shown elsewhere^{10,11} that the second term of Equation (47) describes the dissipative processes of the system while the first term characterizes the “perfect fluid” (e.g., Euler) dynamics. The latter is entirely determined by its functional dependence on the conjugate fields and reduces to the equilibrium pressure tensor of the last section in the case of uniform $\beta(\mathbf{r}, t)$. In the following attention will be restricted to $\overline{t_{0ij}^\ell}(\mathbf{r}|y(t))$ and its possible relationship to an underlying “local equilibrium thermodynamics.”

For the purpose of calculating averages in the rest frame, the macroscopic velocity dependence can be eliminated so the conjugate fields simplify to

$$\{y(\mathbf{r})\} \leftrightarrow \{-\nu(\mathbf{r}), \beta(\mathbf{r}), 0\}. \quad (48)$$

Also, the dependence of these fields on time has been suppressed for simplicity here and below. The hydrodynamic (mechanical) pressure and pressure tensor are defined in terms of the momentum flux as in the previous section

$$p_{ij}^\ell(\mathbf{r}|\beta, \nu) \equiv \overline{t_{0ij}^\ell}(\mathbf{r}|\beta, \nu), \quad (49)$$

$$p_m^\ell(\mathbf{r}|\beta, \nu) \equiv \frac{1}{3} \overline{t_{0ii}^\ell}(\mathbf{r}|\beta, \nu). \quad (50)$$

Since the functional form of the operator $t_{ij}(\mathbf{r}, t)$ in Equation (52) is the same for both equilibrium and nonequilibrium states, the local pressure here is still given by Equation (27) with only the definition of the average changed

$$p_m^\ell(\mathbf{r}|\beta, \nu) = \frac{1}{3} \left(\overline{2K(\mathbf{r})}^\ell + \overline{\mathcal{V}(\mathbf{r})}^\ell \right). \quad (51)$$

Next, a thermodynamics associated with the local equilibrium state is defined in analogy to strict equilibrium via the normalization function $Q^\ell[y]$ of Equation (44). As noted above the average velocity field can be transformed to zero so the energy $e(\mathbf{r})$ is also in the local rest frame. Then Equation (44) becomes

$$Q^\ell[\beta, \nu] = \ln \sum_N Tr^{(N)} e^{-\int d\mathbf{r} (\beta(\mathbf{r}) e(\mathbf{r}) - \nu(\mathbf{r}) n(\mathbf{r}))}. \quad (52)$$

In analogy to Equations (7) and (8), a local equilibrium pressure can be defined by

$$\int d\mathbf{r} \beta(\mathbf{r}) p^\ell(\mathbf{r}|\beta, \nu) \equiv Q^\ell[\beta, \nu], \quad (53)$$

or for extensive systems

$$\frac{1}{V} \int d\mathbf{r} \beta(\mathbf{r}) p^\ell(\mathbf{r}|\beta, \nu) = \frac{\partial Q^\ell[\beta, \nu]}{\partial V} \bigg|_{\beta, \nu}. \quad (54)$$

Carrying out the volume derivative leads to

$$\begin{aligned} \frac{\partial Q^\ell[\beta, \nu]}{\partial V} \bigg|_{\beta, \nu} &= \frac{1}{3V} \sum_{\alpha=1}^N \frac{1}{2m} \left[\overline{p_{\alpha j}^2, \beta(\mathbf{q}_\alpha)} \right]_+ \bigg|_{y(t)} \\ &+ \frac{1}{2} \sum_{\alpha \neq \gamma=1}^N \beta(\mathbf{q}_\alpha) (\mathbf{q}_\gamma - \mathbf{q}_\alpha) \cdot \mathbf{F}_{\alpha\gamma}(|\mathbf{q}_\alpha - \mathbf{q}_\gamma|) \bigg|_{y(t)}, \end{aligned} \quad (55)$$

$$= \frac{1}{V} \int d\mathbf{r} \beta(\mathbf{r}) \frac{1}{3} \left[\overline{2K_0(\mathbf{r})}^\ell + \overline{\mathcal{V}_0(\mathbf{r})}^\ell \right], \quad (56)$$

where $K_0(\mathbf{r})$ and $\mathcal{V}_0(\mathbf{r})$ are given by Equations (14) and (15). Therefore, Equation (54), or equivalently Equation (53) gives the identification

$$Q^\ell[\beta, \nu] = \int d\mathbf{r} \beta(\mathbf{r}) p_0^\ell(\mathbf{r}|\beta, \nu), \quad (57)$$

with

$$p_0^\ell(\mathbf{r}|\beta, \nu) = \frac{1}{3} \left[\overline{2K_0(\mathbf{r})}^\ell + \overline{\mathcal{V}_0(\mathbf{r})}^\ell \right]. \quad (58)$$

This is not the same as the hydrodynamic pressure of Equation (53), $p^\ell(\mathbf{r}, \beta|\nu)$. Their volume integrals are the same

$$P^\ell[\beta, \nu] \equiv \frac{1}{V} \int d\mathbf{r} p_0^\ell(\mathbf{r}|\beta, \nu) = \frac{1}{V} \int d\mathbf{r} p_m^\ell(\mathbf{r}|\beta, \nu), \quad (59)$$

but only $p_0^\ell(\mathbf{r}|\beta, \nu)$ provides the local density for the grand potential

$$Q^\ell[\beta, \nu] = \int d\mathbf{r} \beta(\mathbf{r}) p_0^\ell(\mathbf{r}|\beta, \nu) \neq \int d\mathbf{r} \beta(\mathbf{r}) p_m^\ell(\mathbf{r}|\beta, \nu). \quad (60)$$

This can be stated in an equivalent way. Using Equations (29) and (30) the relationship is

$$p_m^\ell(\mathbf{r}|\beta, \nu) = p_0^\ell(\mathbf{r}|\beta, \nu) + \Delta p_0^\ell(\mathbf{r}|\beta, \nu). \quad (61)$$

$$\partial_j \overline{t_{0ij}^\ell}(\mathbf{r}) = \partial_j \overline{t_{0ji}^\ell}(\mathbf{r}). \quad (68)$$

$$\Delta p_0^\ell(\mathbf{r}|\beta, \nu) = \frac{\hbar^2}{12m} \nabla^2 \bar{n}^\ell(\mathbf{r}) + \frac{1}{6} \sum_{\alpha \neq \sigma=1}^N F_{\alpha\sigma i}(|\mathbf{q}_\alpha - \mathbf{q}_\sigma|) [\mathcal{D}_i(\mathbf{r}, \mathbf{q}_\alpha, \mathbf{q}_\sigma) - (\mathbf{q}_\sigma - \mathbf{q}_\alpha)_i \delta(\mathbf{r} - \mathbf{q}_\alpha)]. \quad (62)$$

Planck's constant has been restored in Equation (62) to make explicit the fact that this term has a purely quantum origin. The local pressures differ for strongly inhomogeneous states, but are the same globally

$$\int d\mathbf{r} \Delta p_0^\ell(\mathbf{r}|\beta, \nu) = 0. \quad (63)$$

However, in contrast to the strict equilibrium case of the last section they do not both give the grand potential due to the appearance of $\beta(\mathbf{q}_\alpha)$ in Equation (55)

$$\int d\mathbf{r} \beta(\mathbf{r}) \Delta p_0^\ell(\mathbf{r}|\beta, \nu) \neq 0. \quad (64)$$

Thus the hydrodynamic local pressure, $p_m^\ell(\mathbf{r}|\beta, \nu)$, does not have the expected relationship to thermodynamics.

The seeming paradox now is that the stationary solution to the hydrodynamic equations is Equation (35)

$$\partial_j p_{ij}^\ell(\mathbf{r}, \beta|\nu) = -\langle n(\mathbf{r}) \rangle \partial_i v^{\text{ext}}(\mathbf{r}), \quad (65)$$

$$p_{ij}^\ell(\mathbf{r}, \beta|\nu) = \frac{1}{3} p_m^\ell(\mathbf{r}, \beta|\nu) \delta_{ij} + \tilde{p}_{ij}^\ell(\mathbf{r}, \beta|\nu), \quad (66)$$

but the corresponding pressure $p_m^\ell(\mathbf{r}, \beta|\nu)$ is not the thermodynamic local pressure $p_0^\ell(\mathbf{r}|\beta, \nu)$ even in the limit of uniform β . The hydrodynamic stationary state is not the thermodynamic state.

4 | REVISED MOMENTUM FLUX AND HYDRODYNAMIC PRESSURE

While there is no flexibility in the choice $p_0^\ell(\mathbf{r}|\beta, \nu)$ in the first equality of Equation (60) due to the space dependence of $\beta(\mathbf{r})$ there is some ambiguity in $p_m^\ell(\mathbf{r}|\beta, \nu)$ available to remove the inequality. This is due to the fact that only the space derivative of $t_{0ij}(\mathbf{r})$ occurs in Equation (40). Consequently, any tensor of the form

$$\overline{t_{0ij}^\ell}(\mathbf{r}) = \overline{t_{0ij}^\ell}(\mathbf{r}) + \varepsilon_{jkn} \partial_k A_{ni}(\mathbf{r}), \quad (67)$$

will give an equivalent derivative. More generally, it is shown that the contour in the definition of $\mathcal{D}_j(\mathbf{r}, \mathbf{q}_1, \mathbf{q}_2)$, Equation (21) can be chosen such that $t_{0ij}(\mathbf{r}) = t_{0ji}(\mathbf{r})$ (see Supporting Information S2), and hence

Then the generalization of Equation (67) to its symmetric form is

$$\overline{t_{0ij}^\ell} = \overline{t_{0ji}^\ell} + \varepsilon_{ikl} \varepsilon_{jmn} \partial_k \partial_m A_{ln}, \quad A_{ln} = A_{nl}, \quad (69)$$

where here $\overline{t_{0ij}^\ell}$ is the symmetric form of Supporting Information S2. It is readily verified that

$$\partial_j \overline{t_{0ij}^\ell}(\mathbf{r}) = \partial_j \overline{t_{0ji}^\ell}(\mathbf{r}), \quad \partial_i \overline{t_{0ij}^\ell}(\mathbf{r}) = \partial_i \overline{t_{0ji}^\ell}(\mathbf{r}). \quad (70)$$

The hydrodynamic pressure tensor associated with $\overline{t_{0ij}^\ell}(\mathbf{r}, t)$ is

$$p'_{ij}(\mathbf{r}|\beta, \nu) \equiv \overline{t_{0ij}^\ell}(\mathbf{r}|\beta, \nu) = \overline{t_{0ji}^\ell}(\mathbf{r}|\beta, \nu) + \varepsilon_{ikl} \varepsilon_{jmn} \partial_k \partial_m A_{ln}(\mathbf{r}|\beta, \nu) = p_{ij}^\ell(\mathbf{r}|\beta, \nu) + \varepsilon_{ikl} \varepsilon_{jmn} \partial_k \partial_m A_{ln}(\mathbf{r}|\beta, \nu), \quad (71)$$

and the corresponding pressure is

$$p'^\ell(\mathbf{r}|\beta, \nu) = p_m^\ell(\mathbf{r}|\beta, \nu) + \frac{1}{3} \varepsilon_{ikl} \varepsilon_{imn} \partial_k \partial_m A_{ln}(\mathbf{r}|\beta, \nu) = p_m^\ell(\mathbf{r}|\beta, \nu) + \frac{1}{3} \left(\partial_k^2 A_{\ell\ell}(\mathbf{r}|\beta, \nu) - \partial_k \partial_\ell A_{\ell k}(\mathbf{r}|\beta, \nu) \right). \quad (72)$$

Use has been made of the identity

$$\varepsilon_{ikl} \varepsilon_{imn} = \delta_{km} \delta_{ln} - \delta_{kn} \delta_{lm}. \quad (73)$$

To further simplify Equation (72) choose the arbitrary tensor $A_{\ell k}(\mathbf{r}|\beta, \nu)$ to be diagonal

$$A_{lk}(\mathbf{r}|\beta, \nu) \rightarrow \delta_{kl} A(\mathbf{r}|\beta, \nu), \quad (74)$$

giving

$$p'^\ell(\mathbf{r}|\beta, \nu) = p^\ell(\mathbf{r}|\beta, \nu) + \frac{2}{3} \nabla^2 A(\mathbf{r}|\beta, \nu), \quad (75)$$

$$p'_{ij}(\mathbf{r}|\beta, \nu) = p_{ij}^\ell(\mathbf{r}|\beta, \nu) + (\delta_{ij} \nabla^2 - \partial_j \partial_i) A(\mathbf{r}|\beta, \nu). \quad (76)$$

It is now seen that the hydrodynamic pressure $p'^\ell(\mathbf{r}|\beta, \nu)$ can be equated to the thermodynamic pressure $p_0^\ell(\mathbf{r}|\beta, \nu)$ by the choice

$$\frac{2}{3} \nabla^2 A(\mathbf{r}|\beta, \nu) = -\Delta p_0^\ell(\mathbf{r}|\beta, \nu). \quad (77)$$

Since $\Delta p_0^\ell(\mathbf{r}|\beta, \nu)$ is given explicitly by Equation (62), this Poisson's equation for $A(\mathbf{r}|\beta, \nu)$ is well-defined. In summary, by modifying the form for the average stress tensor the momentum balance equation is unchanged but the mechanical pressure can be chosen equal to the equilibrium pressure for both the equilibrium and

local equilibrium states, and that pressure is $p_0^e(\mathbf{r}|\beta, \nu)$ or $p_0^e(\mathbf{r}, \beta|\nu)$ respectively.

As a special simple case the explicit results for a noninteracting inhomogeneous gas are

$$\frac{2}{3}\nabla^2 A(\mathbf{r}|\beta, \nu) = -\frac{\hbar^2}{12m}\nabla^2 \bar{n}^e(\mathbf{r}), \quad (78)$$

with the solution

$$A(\mathbf{r}|\beta, \nu) = -\frac{\hbar^2}{8m}\bar{n}^e(\mathbf{r}) + a(\mathbf{r}|\beta, \nu), \quad \nabla^2 a(\mathbf{r}|\beta, \nu) = 0. \quad (79)$$

In summary, by changing the form of the pressure tensor to Equation (71) there is the freedom to choose the new mechanical local pressure $p'^e(\mathbf{r}|\beta, \nu)$ to be equal to the thermodynamic local pressure $p_0^e(\mathbf{r}|\beta, \nu)$. In this way the desired equality

$$Q^e[\beta, \nu] = \int d\mathbf{r} \beta(\mathbf{r}) p_0^e(\mathbf{r}|\beta, \nu) = \int d\mathbf{r} \beta(\mathbf{r}) p'^e(\mathbf{r}|\beta, \nu), \quad (80)$$

is recovered. This result holds for both equilibrium and local equilibrium states.

5 | DISCUSSION

The definition of a local pressure from two conceptually different origins has been considered. The first is thermodynamic in nature, associated with the local density for the grand potential. The other is mechanical in nature, associated with the average of the local momentum flux (referred to as the mechanical or hydrodynamic pressure). First, a strictly equilibrium state for an inhomogeneous system was described. This is the case of interest for density functional theory where the task is to calculate the global free energy. Although local free energy densities are introduced in that context they are mainly for computational convenience. However, they also provide the basis for a local thermodynamics as well. For example, a local pressure follows from a local “Legendre transformation” of the free energy density.⁴ In the present case, the thermodynamic local pressure is introduced directly as the density for the grand potential. It is then compared to the pressure defined from the equilibrium average of the momentum flux (equilibrium force balance equation). The two are different for strongly inhomogeneous states. However, the possibility of adding a contribution $\Delta p_0^e(\mathbf{r}|\beta, \nu)$ whose volume integral vanishes to the grand potential density can be exploited to assure the thermodynamic and hydrodynamic local pressures are the same.

The same analysis for local equilibrium states, where the temperature is also nonuniform, again leads to different forms for the thermodynamic and hydrodynamic local pressures. But in this case there is no longer the flexibility to add a contribution $\Delta p_0^e(\mathbf{r}|\beta, \nu)$ to the thermodynamic local pressure for resolution. Instead, the hydrodynamic

local pressure can be changed by exploiting the fact that the momentum flux occurs in the momentum conservation law only as a divergence of that flux. In this way, agreement of the thermodynamic and hydrodynamic local pressures is restored.

For consistency between the local and strict equilibrium states in the limit of uniform temperature, the local equilibrium form must be used, $p_0^e(\mathbf{r}|\beta, \nu) = p_m^e(\mathbf{r}|\beta, \nu)$, that is, the same average momentum flux should be adopted in each case. This is a somewhat simpler form than the pressure tensor and local pressure than $p_m^e(\mathbf{r}|\beta, \nu)$ for the strict equilibrium case. The requirement that the force balance equation at local equilibrium gives the correct local density for the local equilibrium grand potential gives a strong constraint on the equilibrium form as well.

As noted in Equation (61) both $p_0^e(\mathbf{r}|\beta, \nu)$ and $p_m^e(\mathbf{r}|\beta, \nu)$ are the local densities for the global pressure, without any need for modification of the pressure tensor. If that were chosen to be the constraint of thermodynamic consistency, then no modification of the choice $p_m^e(\mathbf{r}|\beta, \nu) = p^e(\mathbf{r}|\beta, \nu)$ as in the equilibrium case is required—no change in the pressure tensor. Agreement of the equilibrium and local equilibrium cases would be direct, but the inequality Equation (60) would remain. The consequences of this for local equilibrium thermodynamics is not clear and needs to be explored further. The grand potential is a Massieu–Planck functional in the foundations of local equilibrium thermodynamics.¹¹ The pressure functional alone has no corresponding role.

Reference has been made above to density functional theory where the pressure is expressed as a functional of the density n rather than the activity ν . This change of variables is obtained by inverting

$$\overline{n(\mathbf{r}|\beta, \nu)}^e = \bar{n} \rightarrow \nu = \nu(\mathbf{r}|\beta, \bar{n}). \quad (81)$$

This is a difficult problem, separate from the discussion referring to the definition of $p_0^e(\mathbf{r}|\beta, \nu)$. However, the proof that the hydrodynamic pressure can be chosen to be the same as the thermodynamic pressure assures that the tools of density functional theory can be used within the hydrodynamic context as well.

In closing it is useful to return to the extensive literature mentioned above regarding the definition of the pressure tensor and its measurement by molecular dynamics simulation,^{9,23–25} and its common implementation in the Sandia National Laboratories code LAMMPS (Large-scale Atomic/Molecular Massively Parallel Simulator). These simulations refer to methods for direct evaluation of the microscopic definitions of the various components for the pressure tensor. Here, no consideration is given for the pressure tensor components beyond its scalar trace. In that respect the equivalence constraint does not determine the full pressure tensor. Also, only its local equilibrium average is involved—the residual irreversible component of the momentum flux is not affected. A direct measurement of the momentum flux by simulation would give the total of both components (see Equation (47)), which would give the part studied here only for non-dissipative flows, for example, the equilibrium state. Finally, the equivalence condition is the equality of two functionals of the density and temperature. The numerical confirmation of functional equivalence is

indeed a formidable task. Thus, the important simulation studies are only of indirect bearing on the limited scope considered here.

ACKNOWLEDGMENTS

The authors are indebted to S.B. Trickey for comments and criticism of early drafts. This research was supported by US DOE Grant DE-SC0002139. Kai Luo was also supported by U.S. NSF CSEDI grant EAR-1901813 and the Carnegie Institution for Science.

ORCID

James Dufty  <https://orcid.org/0000-0002-2145-3733>

Jeffrey Wrighton  <https://orcid.org/0000-0003-4696-5732>

Kai Luo  <https://orcid.org/0000-0002-3802-0029>

REFERENCES

1. Huang K. *Statistical Mechanics*. New York: John Wiley & Sons; 1987.
2. Evans R. Density functionals in the theory of nonuniform fluids. In: Henderson D, ed. *Fundamentals of Inhomogeneous Fluids*. New York: Marcel Dekker; 1992:85-176.
3. Percus JK. *The Liquid State of Matter: Fluids, Simple and Complex*. Amsterdam, The Netherlands: North-Holland; 1982.
4. Pozhar LA, Gubbins KE, Percus JK. Generalized compressibility equation for inhomogeneous fluids at equilibrium. *Phys Rev E*. 1993;48:1819-1822.
5. Percus JK. The pressure tensor in a non-uniform fluid. *Chem Phys Lett*. 1986;123:311-314.
6. Irving JH, Kirkwood JG. The statistical mechanical theory of transport processes. IV. The equations of hydrodynamics. *J Chem Phys*. 1950;18:817-829.
7. Harasima A. Molecular theory of surface tension. In: Prigogine I, Debye D, eds. *Advances in Chemical Physics*. New York: John Wiley & Sons; 1958:203-237.
8. Schofield P, Henderson JR. Statistical mechanics of inhomogeneous fluids. *Proc R Soc Lond A*. 1982;379:231-246.
9. Walton JPRB, Tildesley DJ, Rowlinson JS, Henderson JR. The pressure tensor at the planar surface of a liquid. *Mol Phys*. 1983;48:1357-1368.
10. Dufty JW, Luo K, Wrighton JM. Generalized hydrodynamics revisited. *Phys Rev Res*. 2020;2:023036.
11. Zubarev D, Ropke G, Morozov V. *Statistical Mechanics of Non-equilibrium Processes*. Akademie Verlag, Berlin; 1996.
12. Hafskjold B, Ikeshoji T. Microscopic pressure tensor for hard-sphere fluids. *Phys Rev E*. 2002;66:011203.
13. Shi K, Shen Y, Santiso EE, Gubbins KE. Microscopic pressure tensor in cylindrical geometry: pressure of water in a carbon nanotube. *J Chem Theory Comput*. 2020;16:5548-5561.
14. Lovett R, Baus M. A molecular theory of the Laplace relation and of the local forces in a curved interface. *J Chem Phys*. 1997;106:635-644.
15. Jaynes ET. Information theory and statistical mechanics. *Phys Rev*. 1957;106:620-630.
16. Jaynes ET. Information theory and statistical mechanics. II. *Phys Rev*. 1957;108:171-190.
17. Robertson HS. *Statistical Thermophysics*. Englewood Cliffs, NJ: Prentice-Hall; 1993.
18. Ván P. Nonequilibrium thermodynamics: emergent and fundamental. *Phil Trans R Soc A*. 2020;378:20200066.
19. Jou D. Relationships between rational extended thermodynamics and extended irreversible thermodynamics. *Phil Trans R Soc A*. 2020;378:20190172.
20. Ter Haar D. *Elements of Statistical Mechanics*. Elsevier, Oxford, UK; 1995.
21. Dufty JW, Trickey SB. Finite temperature scaling in density functional theory. *Mol Phys*. 2016;114:988-996.
22. McLennan JA. *Introduction to Nonequilibrium Statistical Mechanics*. Englewood Cliffs, NJ: Prentice-Hall; 1989.
23. Hardy RJ. Formulas for determining local properties in molecular dynamics simulations: shock waves. *J Chem Phys*. 1982;76:622-628.
24. Todd BD, Evans DJ, Daivis PJ. Pressure tensor for inhomogeneous fluids. *Phys Rev E*. 1995;52:1627-1638.
25. Heyes DM, Smith ER, Dini D, Zaki TA. The equivalence between volume averaging and method of planes definitions of the pressure tensor at a plane. *J Chem Phys*. 2011;135:024512.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Dufty J, Wrighton J, Luo K. Local pressure for inhomogeneous fluids. *AIChE J*. 2021;67(10):e17323. <https://doi.org/10.1002/aic.17323>